

Dilution Robustness for Mean Field Ferromagnets

Adriano Barra^{*†}, Federico Camboni^{*}, Pierluigi Contucci[†]

^{*} Dipartimento di Fisica, Sapienza Università di Roma

[†] Dipartimento di Matematica, Università di Bologna

Abstract

In this work we compare two different random dilutions on a mean field ferromagnet: the first model is built on a Bernoulli-diluted graph while the second lives on a Poisson-diluted one. While it is known that the two models have, in the thermodynamic limit, the same free energy, we investigate on the structural constraints that the two models must fulfill. We rigorously derive for each model the set of identities for the multi-overlaps distribution, using different methods for the two dilutions: constraints in the former model are obtained by studying the consequences of the self-averaging of the internal energy density, while in the latter are obtained by a stochastic-stability technique. Finally we prove that the identities emerging in the two models are the same, showing *robustness* of the ferromagnetic properties of diluted graphs with respect to the details of dilution.

Keywords: diluted graphs, spin glasses, polynomial identities.

1 Introduction

In the past decade an increasing interest has been shown in statistical mechanics built on diluted graphs (see i.e. [3][10][12][27][28][30]). For diluted spin glasses [24][7] this interest is, at least, double motivated: despite their mean field nature, they share with finite-dimensional models the fact that each spin interact with a finite number of other spins. Moreover they are mathematically

equivalent to some random optimization problems (i.e. K-SAT or X-OR-SAT depending on the size of the instantaneous interaction [25][26]).

Although the cases of simpler ferromagnetic models [4][5][19] are not interesting from the point of view of the hard satisfiability interpretation, they are still interesting for their finite connectivity nature and for testing different variations on the topology of the graph they are based on.

With this aim we consider two different ways of diluting the graph [13]: the first ferromagnet has its links distributed according to a Bernoulli probability distribution, the second according to a Poisson.

For these models we compared the properties of a family of linear constraints for the order parameters (known as Aizenman-Contucci polynomials [2][6] in the case of spin glasses).

These relations were investigated earlier [2][21] in the spin glass framework, where they were obtained as a result of the stability of the quenched measure with respect to random perturbation, or equivalently through the bound on the fluctuation of the internal energy. Here we propose them as a test for robustness under dilution.

The methods to approach the identities for the two models are structurally different. For the Poisson case, in fact, the additive law of Poissonian random variables makes possible the direct exploitation of the stochastic stability property. The same strategy is not available for the Bernoulli random variables but, for those, we derive the set of identities from the general bound on the quenched fluctuations (even though, for the sake of completeness, in the Appendix we derive the constraints within this general framework for the Poisson case too). The methods we use are generalizations of those appearing in [8][11][16][17][23][29][15][20][6][1].

Our main result is a rigorous proof of the identities and, especially, the fact that they coincide for the two dilutions.

2 The mean field diluted ferromagnet

We introduce a large number N of sites, labeled by the index $i = 1, \dots, N$, and associate to each of them an Ising variable $\sigma_i = \pm 1$.

We introduce furthermore two families of discrete independent random vari-

ables $\{i_\nu\}, \{j_\nu\}$, uniformly distributed on $1, 2, \dots, N$.

The Hamiltonian $H_N(\sigma)$ of the diluted ferromagnet is expressed through

$$H_N(\sigma) = - \sum_{\nu=1}^x \sigma_{i_\nu} \sigma_{j_\nu} \quad (1)$$

where x does depend on the dilution probability distribution.

For the Bernoullian dilution case the variable x is called k , and defined by

$$E_B[\cdot] = \sum_{k=0}^M \frac{M!}{(M-k)!k!} \left(\frac{\alpha}{N}\right)^k \left(1 - \frac{\alpha}{N}\right)^{M-k} [\cdot], \quad (2)$$

$M = N(N-1)/2$ being the maximum amount of couples $\sigma_i \sigma_j$ existing in the model and α/N the probability that two spins interact.

$\alpha > 0$ plays the role of the connectivity.

The mean and the variance of k are obtained as

$$E_B[k] = \frac{M\alpha}{N}, \quad (3)$$

$$E_B[k^2] - E_B^2[k] = \frac{M\alpha}{N} \left(1 - \frac{\alpha}{N}\right). \quad (4)$$

We will make use in the sequel of following properties of a Bernoulli distribution:

$$E_B[kg(k)] = \frac{M\alpha}{N} E_B[g(k+1)], \quad (5)$$

$$E_B[k^2g(k)] = \frac{M(M-1)\alpha^2}{N^2} E_B[g(k+2)] - \frac{M\alpha}{N} E_B[g(k+1)], \quad (6)$$

$$\frac{d}{d\alpha} E_B[g(k)] = \frac{M}{N} E_B[g(k+1) - g(k)]. \quad (7)$$

For the Poissonian dilution case, we denote x as $\xi_{\alpha N}$, which is a random variable of mean αN , for some $\alpha > 0$ (again defining the connectivity of the model), such that

$$P(\xi_{\alpha N} = k) = \pi(k, \alpha N) = \exp\left(-\alpha N\right) \frac{(\alpha N)^k}{k!}, \quad k = 0, 1, 2, \dots \quad (8)$$

Furthermore, we stress that the Poisson distribution obeys the following properties

$$k\pi(k, \lambda) = \lambda\pi(k-1, \lambda), \quad (9)$$

$$\frac{d}{d\lambda}\pi(k, \lambda) = -\pi(k, \lambda) + \pi(k-1, \lambda)(1 - \delta_{k,0}). \quad (10)$$

As for the Bernoulli case, the average with respect to the Poisson measure will be denoted by the index

$$E_P = \sum_{k=0}^{\infty} \frac{e^{\alpha N} (\alpha N)^k}{k!}. \quad (11)$$

The expectation with respect to all the quenched variables will be denoted by \mathbf{E} and represents the product of the expectation over the dilution distribution and the expectation over the uniformly distributed variables

$$\mathbf{E} = E_{B,P} \cdot \frac{1}{N^2} \sum_{i,j}^{1,N}.$$

The thermodynamic objects we deal with are the partition function

$$Z_N(\alpha, \beta) = \sum_{\{\sigma\}} e^{-\beta H_N(\alpha)}, \quad (12)$$

the quenched intensive free energy

$$A_N(\alpha, \beta) = \frac{1}{N} \mathbf{E} \ln Z_N(\alpha, \beta), \quad (13)$$

the Boltzmann state

$$\omega(g(\sigma)) = \frac{1}{Z_N(\alpha, \beta)} \sum_{\{\sigma_N\}} g(\sigma) e^{-\beta H_N(\alpha)}, \quad (14)$$

the replicated Boltzmann state

$$\Omega(g(\sigma)) = \prod_s \omega^{(s)}(g(\sigma^{(s)})) \quad (15)$$

and the global average $\langle g(\sigma) \rangle$ defined as

$$\langle g(\sigma) \rangle = \mathbf{E}[\Omega(g(\sigma))]. \quad (16)$$

The functional order parameter of the theory is the infinite family of multi-overlaps, defined as

$$q_{1\dots n} = \frac{1}{N} \sum_{i=1}^N \sigma_i^{(1)} \cdots \sigma_i^{(n)},$$

where particular emphasis is due to the magnetization $m = q_1 = (1/N) \sum_{i=1}^N \sigma_i$ and to the two replica overlap $q_{12} = (1/N) \sum_{i=1}^N \sigma_i^1 \sigma_i^2$.

3 Bernoullian diluted case

Identities in the Bernoullian model will be obtained as a consequence of the internal energy self-average [14][20]; before focusing on this procedure, let us recall that

Definition 1 *A quantity $A(\sigma)$ is called self-averaging if*

$$\lim_{N \rightarrow \infty} \langle \left(A(\sigma) - \langle A(\sigma) \rangle \right)^2 \rangle = \lim_{N \rightarrow \infty} \left(\langle A^2(\sigma) \rangle - \langle A(\sigma) \rangle^2 \right) = 0, \quad (17)$$

by which we can remind the

Proposition 1 *Given two regular functions $A(\sigma)$ and $B(\sigma)$, if at least one of them is self-averaging, then the following relation holds*

$$\lim_{N \rightarrow \infty} \langle A(\sigma) B(\sigma) \rangle = \lim_{N \rightarrow \infty} \langle A(\sigma) \rangle \lim_{N \rightarrow \infty} \langle B(\sigma) \rangle \quad (18)$$

Proof

The proof is straightforward. Let us suppose the self-averaging quantity is $B(\sigma)$ and use $A(\sigma)$ as a trial function. Then we have

$$\begin{aligned} 0 &\leq |\langle A(\sigma) B(\sigma) \rangle - \langle A(\sigma) \rangle \langle B(\sigma) \rangle| \\ &= |\langle A(\sigma) B(\sigma) - A(\sigma) \langle B(\sigma) \rangle + \langle A(\sigma) \rangle B(\sigma) - \langle A(\sigma) \rangle \langle B(\sigma) \rangle| \\ &\leq \sqrt{\langle A^2(\sigma) \rangle} \sqrt{\langle (B(\sigma) - \langle B(\sigma) \rangle)^2 \rangle}, \end{aligned} \quad (19)$$

where, in the last passage, we used Cauchy-Schwartz relation. In the thermodynamic the proof becomes completed. \square

The scheme to follow is then clear: using the above proposition as the underlying backbone in the derivation of the constraints in this section, we must, at first, show that the internal energy density of the model self-averages. Then we use as trial functions suitably chosen quantities of the order parameters. The identities follow by evaluating explicitly both the terms of eq. (18): this operation produces several contributions, all involving the order parameters, among which cancelations occur and the remaining part gives the identities.

3.1 Self-averaging of the internal energy density

Once defined $h_l = H(\sigma^{(l)})/N$ as the density of the Hamiltonian evaluated on the generic l^{th} replica and

$$\theta = \tanh(\beta) \quad (20)$$

$$\alpha' = M\alpha/N^2 \xrightarrow{N \rightarrow \infty} \alpha/2 \quad (21)$$

for simplicity, we start with the following

Theorem 1 *In the thermodynamic limit, and in β -average, the internal energy density self-averages*

$$\lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} \mathbb{E} \left(\Omega(h^2) - \Omega(h)^2 \right) d\beta = 0. \quad (22)$$

This Theorem has already been proved in [18] by using essentially the existence of the thermodynamic limit for the free energy density. To make the paper self contained we provide an alternative proof in the appendix.

We can now introduce the following lemma.

Lemma 1 *Let us consider for simplicity the quantity*

$$\Delta G = \sum_{l=1}^s \left[E \left(\Omega(h_l G) - \Omega(h_l) \Omega(G) \right) \right]. \quad (23)$$

For every regular, smooth function G , in β -average, we have

$$\lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} |\Delta G| d\beta = 0. \quad (24)$$

Proof

$$\int_{\beta_1}^{\beta_2} |\Delta G| d\beta \leq \int_{\beta_1}^{\beta_2} \sum_{l=1}^s |\mathbf{E}[\Omega(h_l G) - \Omega(h_l)\Omega(G)]| d\beta \quad (25)$$

$$\leq \int_{\beta_1}^{\beta_2} \sum_{l=1}^s \sqrt{\mathbf{E}[(\Omega(h_l G) - \Omega(h_l)\Omega(G))^2]} d\beta \quad (26)$$

$$\leq s \int_{\beta_1}^{\beta_2} \sqrt{\mathbf{E}[\Omega(h^2) - \Omega^2(h)]} d\beta \quad (27)$$

$$\leq s \sqrt{\beta_2 - \beta_1} \sqrt{\int_{\beta_1}^{\beta_2} \mathbf{E}[\Omega(h^2) - \Omega^2(h)] d\beta} \xrightarrow{N \rightarrow \infty} 0 \quad (28)$$

where (25) comes from triangular inequality; (26) is obtained via the Jensen inequality applied to the measure $\mathbf{E}[\cdot]$. In the same way (27) comes from Schwarz inequality applied on the measure $\Omega(\cdot)$ (being G bounded), while (28) is obtained via Jensen inequality applied on the measure $(\beta_2 - \beta_1)^{-1} \int_{\beta_1}^{\beta_2} (\cdot) d\beta$. \square

Now we can state the main theorem for the linear constraints.

We are going to introduce a specific trial functions that we call $f_G(\alpha, \beta)$.

Theorem 2 *Let us consider the following series of multi-overlap functions G*

acting, in complete generality, on s replicas

$$\begin{aligned}
f_G(\alpha, \beta) = & \alpha' \left[\left(\sum_{l=1}^s \langle Gm_l^2 \rangle - s \langle Gm_{s+1}^2 \rangle \right) (1 - \theta^2) + \right. \\
& + 2\theta \left(\sum_{a < l}^{1,s} \langle Gq_{al}^2 \rangle - s \sum_l^{1,s} \langle Gq_{l,s+1}^2 \rangle + \frac{s(s+1)}{2} \langle Gq_{s+1,s+2}^2 \rangle \right) + \\
& + 3\theta^2 \left(\sum_{l < a < b}^{1,s} \langle Gq_{l,a,b}^2 \rangle - s \sum_{l < a}^{1,s} \langle Gq_{l,a,s+1}^2 \rangle + \frac{s(s+1)}{2} \sum_l^{1,s} \langle Gq_{l,s+1,s+2}^2 \rangle + \right. \\
& \left. \left. - \frac{s(s+1)(s+2)}{3!} \langle Gq_{s+1,s+2,s+3}^2 \rangle \right) + O(\theta^3) \right], \tag{29}
\end{aligned}$$

in the thermodynamic limit the following generator of linear constraints holds:

$$\lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} d\beta |f_G(\alpha, \beta)| = 0. \tag{30}$$

Proof

Let us consider explicitly the quantities encoded in (23). For the sake of clearness all the calculations are reported in appendix, here we present just the results.

$$\begin{aligned}
\mathbf{E}[\Omega(h_l G)] = & -\alpha' \left[\langle Gm_l^2 \rangle + \theta \left(\sum_{a=1}^s \langle Gq_{a,l}^2 \rangle - s \langle Gq_{l,s+1}^2 \rangle \right) + \right. \\
& + \theta^2 \left(\sum_{a < b}^{1,s} \langle Gq_{l,a,b}^2 \rangle - s \sum_a^{1,s} \langle Gq_{l,a,s+1}^2 \rangle + \frac{s(s+1)}{2} \langle Gq_{l,s+1,s+2}^2 \rangle \right) \\
& \left. + O(\theta^2) \right], \tag{31}
\end{aligned}$$

$$\begin{aligned}
\mathbf{E}[\Omega(h_l)\Omega(G)] &= -\alpha' \left[\langle Gm_l^2 \rangle + \theta \left(\sum_{a=1}^{s+1} \langle Gq_{a,l}^2 \rangle - (s+1) \langle Gq_{l,s+1}^2 \rangle \right) + \right. \\
&\quad + \theta^2 \left(\sum_a^{1,s} \langle Gm_a^2 \rangle - (s+1) \langle Gm_l^2 \rangle + \sum_{a < b}^{1,s} \langle Gq_{l,a,b}^2 \rangle + \right. \\
&\quad \left. - (s+1) \sum_a^{1,s} \langle Gq_{l,a,s+1}^2 \rangle + \frac{(s+1)(s+2)}{2} \langle Gq_{l,s+1,s+2}^2 \rangle \right) \\
&\quad \left. + O(\theta^2) \right]. \tag{32}
\end{aligned}$$

Subtracting the last equation from the former, immediately we conclude that

$$\Delta G = -f_G(\alpha, \beta), \tag{33}$$

from which theorem thesis follows. \square

3.2 Linear constraints for multi-overlaps

We outline here the first order identities as it is customary to do in the spin-glasses counterpart [6] or in neural network [9].

Proposition 2 *The first class of multi-overlap constraints is obtained by choosing $G = m^2$.*

In fact, if we set $G = q_1^2 = m^2$, the function $f_G(\alpha, \beta)$ becomes

$$\begin{aligned}
f_{m^2}(\alpha, \beta) &= \alpha' \left[\left(\langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle \right) (1 - \theta^2) + \right. \\
&\quad \left. - 2\theta \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle \right) + \right. \\
&\quad \left. + 3\theta^2 \left(\langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle \right) + O(\theta^3) \right],
\end{aligned}$$

from which, changing the Jacobian $d\theta = (1 - \theta^2)d\beta$, we get

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} |f_{m^2}(\alpha, \beta)| d\beta &= \frac{\alpha}{2} \int_{\theta_1}^{\theta_2} d \frac{\theta}{(1 - \theta^2)} \left[\left(\langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle \right) + \right. \\ &\quad - 2\theta \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle \right) + \\ &\quad + 3\theta^2 \left(\langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle \right) \\ &\quad \left. + O(\theta^3) \right] = 0, \end{aligned} \quad (34)$$

where, the (not interesting) breakdown at $\theta = 1$, of the expression above, reflects the lack of convergence of the harmonic series we used in eq. (62).

Proposition 3 *The second class of multi-overlap constraints is obtained by choosing $G = q_{12}^2$.*

In fact, if we set $G = q_{12}^2$, the function $f_G(\alpha, \beta)$ becomes

$$\begin{aligned} f_{q_{12}^2}(\alpha, \beta) &= \alpha' \left[\left(2\langle m_1^2 q_{12}^2 \rangle - 2\langle m_3^2 q_{12}^2 \rangle \right) (1 - \theta^2) + \right. \\ &\quad + 2\theta \left(\langle q_{12}^4 \rangle - 4\langle q_{12}^2 q_{23}^2 \rangle + 3\langle q_{12}^2 q_{34}^2 \rangle \right) + \\ &\quad \left. - 6\theta^2 \left(\langle q_{12}^2 q_{123}^2 \rangle - 3\langle q_{12}^2 q_{234}^2 \rangle + 2\langle q_{12}^2 q_{345}^2 \rangle \right) + O(\theta^3) \right]. \end{aligned}$$

Again

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} |f_{q^2}(\alpha, \beta)| d\beta &= \frac{\alpha}{2} \int_{\theta_1}^{\theta_2} d \frac{\theta}{(1 - \theta^2)} \left[\left(\langle m_1^2 q_{12}^2 \rangle - \langle m_3^2 q_{12}^2 \rangle \right) + \right. \\ &\quad + \theta \left(\langle q_{12}^4 \rangle - 4\langle q_{12}^2 q_{23}^2 \rangle + 3\langle q_{12}^2 q_{34}^2 \rangle \right) + \\ &\quad - 3\theta^2 \left(\langle q_{12}^2 q_{123}^2 \rangle - 3\langle q_{12}^2 q_{234}^2 \rangle + 2\langle q_{12}^2 q_{345}^2 \rangle \right) \\ &\quad \left. + O(\theta^3) \right] = 0, \end{aligned} \quad (35)$$

from which the constraints are obtained as the r.h.s. of (34,35) set to zero. \square

4 Poissonian diluted case

To tackle the Poisson diluted ferromagnet, we are going to use the cavity field approach.

The idea beyond this technique is that, calling $F(\beta)$ the extensive free energy and $f(\beta)$ the intensive one, a bridge among the two, in the large N limit, is offered simply by the relation

$$\left(-F_{N+1}(\beta) - F_N(\beta) \right) = f(\beta) + O(N^{-1}). \quad (36)$$

As our system has a topologically quenched disordered the $N + 1$ spin can be seen as an "external random cavity field" for the former system of N particles. The identities derived by tuning that field are called of "stochastic stability". The simplest way to find them is to consider monomials which are left invariant by the random field: the derivative with respect to it, being zero, will produce the desired polynomial.

4.1 Cavity field decompositions for the pressure density

To start applying the sketched plan let us decompose (in distribution) a Poissonian random Hamiltonian of $N + 1$ spins in two Hamiltonians [1]: The former of the "inner" N interacting spins, the latter as the remaining spin interacting with the internal N spins of the cavity.

Up to negligible corrections that go to zero in the thermodynamic limit, we can write in distribution

$$H_{N+1}(\alpha) = - \sum_{\nu=1}^{P_{\alpha(N+1)}} \sigma_{i_\nu} \sigma_{j_\nu} \quad \sim \quad - \sum_{\nu=1}^{P_{\tilde{\alpha}N}} \sigma_{i_\nu} \sigma_{j_\nu} - \sum_{\nu=1}^{P_{2\tilde{\alpha}}} \sigma_{i_\nu} \sigma_{N+1}, \quad (37)$$

or simply for compactness

$$H_{N+1}(\alpha) \sim H_N(\tilde{\alpha}) + \hat{H}_N(\tilde{\alpha}) \sigma_{N+1} \quad (38)$$

where

$$\tilde{\alpha} = \frac{N}{N+1} \alpha \xrightarrow{N \rightarrow \infty} \alpha, \quad \hat{H}_N(\tilde{\alpha}) = - \sum_{\nu=1}^{P_{2\tilde{\alpha}}} \sigma_{i_\nu}. \quad (39)$$

It is useful now to introduce an interpolating parameter $t \in [0, 1]$ on the term encoding the linear connectivity shift so to manage the derivative with respect to the random field by differentiating with respect to this parameter.

Definition 2 *We define the t -dependent Boltzmann state $\tilde{\omega}_t$ as*

$$\tilde{\omega}_t(g(\sigma)) = \frac{1}{Z_{N,t}(\alpha, \beta)} \sum_{\{\sigma\}} g(\sigma) e^{\beta \sum_{\nu=1}^{P\tilde{\alpha}N} \sigma_{i_\nu} \sigma_{j_\nu} + \beta \sum_{\nu=1}^{P_2\tilde{\alpha}t} \sigma_{i_\nu}}. \quad (40)$$

We stress the simplicity by which the t parameter switches among the system of $N + 1$ spins and the one built just by the former N , in the large N limit: In fact, being the two body Hamiltonian left invariant by the gauge symmetry $\sigma_i \rightarrow \epsilon \sigma_i$ for all $i \in (1, \dots, N)$ with $\epsilon = \pm 1$, by choosing $\epsilon = \sigma_{N+1}$ we have

$$Z_{N,t=1}(\tilde{\alpha}, \beta) = Z_{N+1}(\alpha, \beta), \quad (41)$$

$$Z_{N,t=0}(\tilde{\alpha}, \beta) = Z_N(\tilde{\alpha}, \beta). \quad (42)$$

Note that $Z_{N,t}(\tilde{\alpha}, \beta)$ is defined accordingly to (40) and coherently, dealing with the perturbed Boltzmann measure, we introduce an index t also to the global averages $\langle \cdot \rangle \rightarrow \langle \cdot \rangle_t$.

4.2 Stochastic stability via cavity fields

We are now ready to attack the problem.

We divide the ensemble of overlap monomials in two large categories: stochastically stable monomials and (as a side results) not stochastically stable ones. Then we find explicitly the family of the stochastically stable monomials, and by putting their t -derivative equal to zero we obtain the identities. To follow the plan let us introduce the

Definition 3 *We define as stochastically stable monomials those multi-overlap monomials where each replica appears an even number of times.*

We are ready to introduce the main theorem, which offers, as a straightforward consequence, a useful corollary, stated immediately after.

Theorem 3 *At $t = 1$ the Boltzmannfaktor of the perturbed measure is comparable with the canonical Boltzmannfaktor, and, in the thermodynamic limit, we get*

$$\lim_{N \rightarrow \infty} \mathbb{E} \tilde{\Omega}_{N,t=1}(\sigma_{i_1} \sigma_{i_2} \dots \sigma_{i_n}) = \lim_{N \rightarrow \infty} \mathbb{E} \Omega_{N+1}(\sigma_{i_1} \sigma_{i_2} \dots \sigma_{i_n} \sigma_{N+1}^n). \quad (43)$$

Corollary 1 *In the thermodynamic limit, the averages $\langle \cdot \rangle_t$ of the stochastically stable monomials become t -independent in β -average.*

Proof

Let us focus on the proof of Theorem 3. Corollary 1 will be produced as a straightforward application of Theorem 3 on stochastically stable monomials. Let us start the proof. Let us assume for a generic multi-overlap monomial the following representation

$$Q = \sum_{i_l^1} \dots \sum_{i_l^s} \prod_{l=1}^{n^a} \sigma_{i_l^a}^a I(\{i_l^a\})$$

where a labels replicas, the inner product accounts for the spins depicted by the index l which belong to the Boltzmann state a of the product state Ω for the multi-overlap $q_{a,a'}$ and runs over the integers from 1 to the amount of times the replica a appears into the expression.

The external product multiplies all the terms coming from the internal one. The factor I fixes replica-bond constraints.

For example the monomial $Q = q_{12} q_{23}$ has $s = 3, n^1 = n^3 = 1, n^2 = 2$ and $I = N^{-2} \delta_{i_1^1, i_1^2} \delta_{i_1^2, i_2^3}$, there the δ -functions give the correlations $1, 2 \rightarrow q_{1,2}$ and $2, 3 \rightarrow q_{2,3}$.

By applying the Boltzmann and quenched-disordered expectations we get

$$\langle Q \rangle_t = \mathbf{E} \sum_{i_l^a} I(\{i_l^a\}) \prod_{a=1}^s \omega_t \left(\prod_{l=1}^{n^a} \sigma_{i_l^a}^a \right).$$

Let us suppose now that Q is not stochastically stable (for otherwise the proof will be simply ended) and let us decompose it by factorizing the Boltzmann state ω and splitting the terms involving replicas appearing an even number

of times from the ones involving replicas appearing an odd number of times. Then, evaluate the whole receipt at $t = 1$.

$$\langle Q \rangle_t = \mathbf{E} \sum_{i_l^a, i_l^b} I(\{i_l^a\}, \{i_l^b\}) \prod_{a=1}^u \omega_a \left(\prod_{l=1}^{n^a} \sigma_{i_l^a}^a \right) \prod_{b=u+1}^s \omega_b \left(\prod_{l=1}^{n^b} \sigma_{i_l^b}^b \right),$$

where u stands for the amount of replicas which appear an odd number of times inside Q .

In this way we split the measure Ω in two ensembles ω_a and ω_b . Replicas belonging to ω_b are an even number while the ones in ω_a an odd number.

At this point, as the Hamiltonian has two body interaction and consequently is left unchanged by the symmetry $\sigma_i^a \rightarrow \sigma_i^a \sigma_{N+1}^a, \forall i \in (1, N)$ (as $\sigma_{N+1}^2 \equiv 1$), we apply such a symmetry globally to the whole set of N spins. The even measure is left unchanged by this symmetry while the odd one takes a multiplying term σ_{N+1}

$$\langle Q \rangle = \sum_{i_l^a, i_l^b} I(\{i_l^a\}, \{i_l^b\}) \prod_{a=1}^u \omega(\sigma_{N+1}^a \prod_{l=1}^{n^a} \sigma_{i_l^a}^a) \prod_{b=u+1}^s \omega(\sigma_{N+1}^b \prod_{l=1}^{n^b} \sigma_{i_l^b}^b).$$

The last trick is that, by noticing the arbitrariness of the $N+1$ label in σ_{N+1} , we can change it to a generic label k for each $k \neq \{i_l^a\}$ and multiply by $1 = N^{-1} \sum_{k=1}^N$. At finite N the thesis is recovered forgetting terms $O(1/N)$ and becomes exact in the thermodynamic limit. \square

It is straightforward to check that the effect of Theorem 3 is not felt by stochastically stable multi-overlap monomials (Corollary 1) thanks to the dichotomy of the Ising spins ($\sigma_{N+1}^{2n} \equiv 1 \forall n \in \mathbf{N}$). \square

The last point missing to obtain the identities is finding a streaming equation to work out the derivatives with respect to the random field of the stochastically stable monomials. To this task we introduce the following

Proposition 4 *Given F_s as a generic function of the spins of s replicas, the*

following streaming equation holds

$$\begin{aligned}
\frac{\partial \langle F_s \rangle_{t, \tilde{\alpha}}}{\partial t} &= 2\tilde{\alpha}\theta \left[\sum_{a=1}^s \langle F_s \sigma_{i_0}^a \rangle_{t, \tilde{\alpha}} - s \langle F_s \sigma_{i_0}^{s+1} \rangle_{t, \tilde{\alpha}} \right] + \\
&+ 2\tilde{\alpha}\theta^2 \left[\sum_{a < b}^{1, s} \langle F_s \sigma_{i_0}^a \sigma_{i_0}^b \rangle_{t, \tilde{\alpha}} - s \sum_{a=1}^s \langle F_s \sigma_{i_0}^a \sigma_{i_0}^{s+1} \rangle_{t, \tilde{\alpha}} + \right. \\
&+ \frac{s(s+1)}{2!} \langle F_s \sigma_{i_0}^{s+1} \sigma_{i_0}^{s+2} \rangle_{t, \tilde{\alpha}} \left. \right] + \\
&+ 2\tilde{\alpha}\theta^3 \left[\sum_{a < b < c}^{1, s} \langle F_s \sigma_{i_0}^a \sigma_{i_0}^b \sigma_{i_0}^c \rangle_{t, \tilde{\alpha}} - s \sum_{a < b}^{1, s} \langle F_s \sigma_{i_0}^a \sigma_{i_0}^b \sigma_{i_0}^{s+1} \rangle_{t, \tilde{\alpha}} + \right. \\
&+ \frac{s(s+1)}{2!} \sum_{a=1}^s \langle F_s \sigma_{i_0}^a \sigma_{i_0}^{s+1} \sigma_{i_0}^{s+2} \rangle_{t, \tilde{\alpha}} + \\
&+ \left. \frac{s(s+1)(s+2)}{3!} \langle F_s \sigma_{i_0}^{s+1} \sigma_{i_0}^{s+2} \sigma_{i_0}^{s+3} \rangle_{t, \tilde{\alpha}} \right] + O(\theta^3) \tag{44}
\end{aligned}$$

Proof

The proof follows by direct calculations

$$\begin{aligned}
\frac{\partial \langle F_s \rangle_{t, \tilde{\alpha}}}{\partial t} &= \frac{\partial}{\partial t} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} F_s e^{\sum_{a=1}^s (\beta \sum_{\nu=1}^{P_{\tilde{\alpha}} N} \sigma_{i_\nu}^a \sigma_{j_\nu}^a + \beta \sum_{\nu=1}^{P_{2\tilde{\alpha}} t} \sigma_{i_\nu}^a)} }{\sum_{\{\sigma\}} e^{\sum_{a=1}^s (\beta \sum_{\nu=1}^{P_{\tilde{\alpha}} N} \sigma_{i_\nu}^a \sigma_{j_\nu}^a + \beta \sum_{\nu=1}^{P_{2\tilde{\alpha}} t} \sigma_{i_\nu}^a)} } \right] = \tag{45} \\
&= 2\tilde{\alpha} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} F_s e^{\sum_{a=1}^s (\beta \sigma_{i_0}^a + \beta \sum_{\nu=1}^{P_{\tilde{\alpha}} N} \sigma_{i_\nu}^a \sigma_{j_\nu}^a + \beta \sum_{\nu=1}^{P_{2\tilde{\alpha}} t} \sigma_{i_\nu}^a)} }{\sum_{\{\sigma\}} e^{\sum_{a=1}^s (\beta \sigma_{i_0}^a + \beta \sum_{\nu=1}^{P_{\tilde{\alpha}} N} \sigma_{i_\nu}^a \sigma_{j_\nu}^a + \beta \sum_{\nu=1}^{P_{2\tilde{\alpha}} t} \sigma_{i_\nu}^a)} } \right] - 2\tilde{\alpha} \langle F_s \rangle_{t, \tilde{\alpha}} = \\
&= 2\tilde{\alpha} \mathbf{E} \left[\frac{\tilde{\Omega}_t(F_s e^{\sum_{a=1}^s \beta \sigma_{i_0}^a})}{\tilde{\Omega}_t(e^{\sum_{a=1}^s \beta \sigma_{i_0}^a})} \right] - 2\tilde{\alpha} \langle F_s \rangle_{t, \tilde{\alpha}} = \\
&= 2\tilde{\alpha} \mathbf{E} \left[\frac{\tilde{\Omega}_t(F_s \Pi_{a=1}^s (\cosh \beta + \sigma_{i_0}^a \sinh \beta))}{\tilde{\Omega}_t(\Pi_{a=1}^s (\cosh \beta + \sigma_{i_0}^a \sinh \beta))} \right] - 2\tilde{\alpha} \langle F_s \rangle_{t, \tilde{\alpha}} = \\
&= 2\tilde{\alpha} \mathbf{E} \left[\frac{\tilde{\Omega}_t(F_s \Pi_{a=1}^s (1 + \sigma_{i_0}^a \theta))}{(1 + \tilde{\omega}_t(\sigma_{i_0}^a) \theta)^s} \right] - 2\tilde{\alpha} \langle F_s \rangle_{t, \tilde{\alpha}},
\end{aligned}$$

then, by noting that

$$\Pi_{a=1}^s(1 + \sigma_{i_0}^a \theta) = 1 + \sum_{a=1}^s \sigma_{i_0}^a \theta + \sum_{a < b}^{1,s} \sigma_{i_0}^a \sigma_{i_0}^b \theta^2 + \sum_{a < b < c}^{1,s} \sigma_{i_0}^a \sigma_{i_0}^b \sigma_{i_0}^c \theta^3 + \dots$$

$$\frac{1}{(1 + \tilde{\omega}_t \theta)^s} = 1 - s \tilde{\omega}_t \theta + \frac{s(s+1)}{2!} \tilde{\omega}_t^2 \theta^2 - \frac{s(s+1)(s+2)}{3!} \tilde{\omega}_t^3 \theta^3 + \dots$$

we get

$$\begin{aligned} \frac{\partial \langle F_s \rangle_{t, \tilde{\alpha}}}{\partial t} &= 2\tilde{\alpha} \mathbf{E}[\tilde{\Omega}_t(F_s(1 + \sum_{a=1}^s \sigma_{i_0}^a \theta + \sum_{a < b}^{1,s} \sigma_{i_0}^a \sigma_{i_0}^b \theta^2 + \sum_{a < b < c}^{1,s} \sigma_{i_0}^a \sigma_{i_0}^b \sigma_{i_0}^c \theta^3 + \dots)) \times \\ &\quad \times (1 - s \tilde{\omega}_t \theta + \frac{s(s+1)}{2!} \tilde{\omega}_t^2 \theta^2 - \frac{s(s+1)(s+2)}{3!} \tilde{\omega}_t^3 \theta^3 + \dots)] - 2\tilde{\alpha} \langle F_s \rangle_{t, \tilde{\alpha}}, \end{aligned}$$

from which the thesis follows. \square

4.3 Linear constraints for multi-overlaps

We saw that the stochastically stable multi-overlap monomials become asymptotically independent by the t parameter upon increasing the size of the system. Calling for simplicity $G_N(q)$ a stochastically stable multi-overlap monomial, identities follow as a consequence of Corollary 1 and are encoded in the following relation

$$\lim_{N \rightarrow \infty} \partial_t \langle G_N(q) \rangle_t = 0.$$

As we did when we investigated the Bernoullian model, we analyze the stability of $\langle m^2 \rangle$ and $\langle q_{12}^2 \rangle$, up to the third order in θ , so to compare the results at the end.

$$\begin{aligned} \partial_t \langle m_1^2 \rangle_t &= 2\tilde{\alpha} \theta \left(\langle m_1^3 \rangle_t - \langle m_1^2 m_2 \rangle_t \right) - 2\tilde{\alpha} \theta^2 \left(\langle m_1^2 q_{12} \rangle_t - \langle m_1^2 q_{23} \rangle_t \right) \\ &\quad + 2\tilde{\alpha} \theta^3 \left(\langle m_1^2 q_{123} \rangle_t - \langle m_1^2 q_{234} \rangle_t \right) + O(\theta^3) \end{aligned}$$

$$\begin{aligned} \Rightarrow \quad & \left[2\alpha\theta \left(\langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle \right) - 2\alpha\theta^2 \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle \right) \right. \\ & \left. + 2\alpha\theta^3 \left(\langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle \right) + O(\theta^3) \right] = 0 \end{aligned} \quad (46)$$

$$\begin{aligned} \partial_t \langle q_{12}^2 \rangle_t &= 4\tilde{\alpha}\theta \left(\langle m_1 q_{12}^2 \rangle_t - \langle m_3 q_{12}^2 \rangle_t \right) + 2\tilde{\alpha}\theta^2 \left(\langle q_{12}^3 \rangle_t - 4\langle q_{12}^2 q_{13} \rangle_t + 3\langle q_{12}^2 q_{34} \rangle_t \right) + \\ &- 4\tilde{\alpha}\theta^3 \left(\langle q_{12}^2 q_{123} \rangle_t - 3\langle q_{12}^2 q_{134} \rangle_t + 4\langle q_{12}^2 q_{345} \rangle_t \right) + O(\theta^3) \\ \Rightarrow \quad & \left[4\alpha\theta \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_3^2 q_{12}^2 \rangle \right) + 2\alpha\theta^2 \left(\langle q_{12}^4 \rangle - 4\langle q_{12}^2 q_{13}^2 \rangle + 3\langle q_{12}^2 q_{34}^2 \rangle \right) + \right. \\ & \left. - 4\alpha\theta^3 \left(\langle q_{12}^2 q_{123}^2 \rangle - 3\langle q_{12}^2 q_{134}^2 \rangle + 2\langle q_{12}^2 q_{345}^2 \rangle \right) + O(\theta^3) \right] = 0 \end{aligned} \quad (47)$$

5 Discussion and outlook

Let us start this section by comparing the results we get from the two models. We have to compare respectively eq.s (34) versus (46) and (35) versus (47). We see that the details of the dilution do not affect the constraints: the series show the same set of identities.

In fact, despite we are not generally allowed to set to zero each term in the expressions (46,47,34,35) (as we do to obtain alone the following identities (48-53)), at least close to the critical line, where different multi-overlaps have different scaling laws [1], i.e. $q_n^2 \propto (\alpha\theta - 1)^n$, such a spreading is possible and we can forget each single (α, β) -coefficient as it does not affect the identities (it is never involved into the averages $\langle \cdot \rangle$).

We get

$$0 = \langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle, \quad (48)$$

$$0 = \langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle, \quad (49)$$

$$0 = \langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle, \quad (50)$$

when investigating the magnetization as a trial function and

$$0 = \langle m_1^2 q_{12}^2 \rangle - \langle m_3^2 q_{12}^2 \rangle, \quad (51)$$

$$0 = \langle q_{12}^4 \rangle - 4\langle q_{12}^2 q_{23}^2 \rangle + 3\langle q_{12}^2 q_{34}^2 \rangle, \quad (52)$$

$$0 = \langle q_{12}^2 q_{123}^2 \rangle - 3\langle q_{12}^2 q_{134}^2 \rangle + 2\langle q_{12}^2 q_{345}^2 \rangle, \quad (53)$$

when investigating the two replica overlap.

Even if a minor point, we stress that the differences among the global (α, β) -coefficients (not shown here) clearly are related to the differences in the two method involved (in the former the constraints appear under the integral over the temperature, while in the latter this β -average is already worked out), furthermore the limiting connectivity in the Bernoulli dilution is $\alpha/2$, while is 2α in the Poisson model; so there is an overall factor 4 of difference among the results (for the sake of clearness we worked out in the appendix also the constraints in the Poisson diluted case via the first method, to check explicitly the coherence among the two model coefficients).

Then, by looking explicitly at the constraints, several physical features can be recognized, in fact every term is well known:

the first class (Eq.s 48,49,50) is the standard first momentum self-averaging on replica symmetric systems; In fact, by assuming replica equivalence, eq. (48) turns out to be the standard internal energy self-averaging of the Curie-Weiss model. Eq.(49) and (50) contribute as higher order internal energy self-average by taking into account the dilution (in fact, they go to zero whenever $\alpha \rightarrow \infty$ because θ -powers higher than one go to zero and only the Curie-Weiss self-averaging for the internal energy survives as it should).

With a glance at the identities coming from the second constraint series (Eq.s 51,52,53) we recognize immediately the replica symmetry ansatz for the magnetization in the first identity, followed by the first and the second Aizenman-Contucci relation for systems with quenched disorder [2][4].

Interestingly these series are in agreement even with other models, apparently quite far away, as spin-glasses with Gaussian coupling $\mathcal{N}[1, 1]$ instead of $\mathcal{N}[0, 1]$ [15]. A very interesting conjecture may be that these constraints hold for systems whose interaction has on average positive strength and are affected by quenched disorder, independently if the disorder affects the strength of the interaction or the topology of the interaction. We plan to report soon on this topics.

Acknowledgment

The author are pleased to thank an anonymous referee for useful suggestions. AB work in this paper is partially supported by the Technological Vaucher Contract n.11606 of Calabria Region and partially by the CULTAPTATION Project (European Commission contract FP6 - 2004-NEST-PATH-043434). PC acknowledge Strategic Grant of University of Bologna.

A Appendix

A.1 Alternative proof of energy self-averaging

Starting from the thermodynamic relation

$$\mathbf{E}[\Omega(h^2) - \Omega^2(h)] = -\frac{1}{N} \frac{d}{d\beta} \mathbf{E}[\Omega(h)] \quad (54)$$

we evaluate explicitly the term $E[\Omega(h)]$ as

$$\mathbf{E}[\Omega(h)] = -\frac{1}{N} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sum_{\nu=1}^k \sigma_{i_\nu} \sigma_{j_\nu} e^{-\beta H}}{Z_N(\alpha, \beta)} \right] = \quad (55)$$

$$= -\frac{1}{N} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} k \sigma_{i_0} \sigma_{j_0} e^{-\beta H}}{Z_N(\alpha, \beta)} \right] = \quad (56)$$

$$= -\alpha' \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0} \sigma_{j_0} e^{\beta \sigma_{i_0} \sigma_{j_0}} e^{-\beta H}}{\sum_{\{\sigma\}} e^{\beta \sigma_{k_0} \sigma_{l_0}} e^{-\beta H}} \right] = \quad (57)$$

$$= -\alpha' \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0} \sigma_{j_0} (\cosh \beta + \sigma_{i_0} \sigma_{j_0} \sinh \beta) e^{-\beta H}}{\sum_{\{\sigma\}} (\cosh \beta + \sigma_{k_0} \sigma_{l_0} \sinh \beta) e^{-\beta H}} \right] = \quad (58)$$

$$= -\alpha' \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0} \sigma_{j_0} (1 + \sigma_{i_0} \sigma_{j_0} \theta) e^{-\beta H}}{\sum_{\{\sigma\}} (1 + \sigma_{k_0} \sigma_{l_0} \theta) e^{-\beta H}} \right] = \quad (59)$$

$$= -\alpha' \mathbf{E} \left[\frac{\omega(\sigma_{i_0} \sigma_{j_0}) + \theta}{1 + \omega(\sigma_{k_0} \sigma_{l_0}) \theta} \right], \quad (60)$$

where in (56) we fixed the index ν , in (57) we used the property (5) of the Bernoulli distribution and we introduced two further families of random variables $\{k_\nu\}, \{l_\nu\}$, and in (58) we used $e^{\beta \sigma_{i_0} \sigma_{j_0}} = \cosh \beta + \sigma_{i_0} \sigma_{j_0} \sinh \beta$.

Let us now expand the denominator of (60) keeping in mind the relation

$$\frac{1}{(1 + \tilde{\omega}_t \theta)^p} = 1 - p\tilde{\omega}_t \theta + \frac{p(p+1)}{2!} \tilde{\omega}_t^2 \theta^2 - \frac{p(p+1)(p+2)}{3!} \tilde{\omega}_t^3 \theta^3 + \dots$$

such that, by posing $p = 1$, we obtain

$$\mathbf{E}[\Omega(h)] = -\alpha' \mathbf{E} \left[\theta + \sum_{n=1}^{\infty} (-1)^n \theta^n (1 - \theta^2) \langle q_{1\dots n}^2 \rangle \right]. \quad (61)$$

By applying the modulus function to the equation above we can proceed further with the following bound

$$|\mathbf{E}[\Omega(h)]| \leq \alpha' \mathbf{E} \left[|\theta| + \sum_{n=1}^{\infty} |\theta^n (1 - \theta^2) \langle q_{1\dots n}^2 \rangle| \right]. \quad (62)$$

Both $|\theta|$ and $|\langle q_{1\dots n}^2 \rangle|$ belong to $[0, 1]$ so we get

$$|E[\Omega(h)]| \leq \alpha' \left[1 + (1 - \theta^2) \sum_{n=1}^{\infty} \theta^n \right], \quad (63)$$

whose harmonic series converges to $1/(1 - \theta)$, $|\theta| < 1$;

The fact that the convergence is not guaranteed at zero temperature with this technique is not a problem because, first the identities we are looking for hold in β -average, secondly the zero temperature has been intensively investigated elsewhere [22].

For each finite β , then, we can write

$$|\mathbf{E}[\Omega(h)]| \leq \alpha' \left[1 + \frac{(1 - \theta^2)}{1 - \theta} \right] = \quad (64)$$

$$= \alpha' \left[1 + \frac{(1 - \theta)(1 + \theta)}{1 - \theta} \right] = \quad (65)$$

$$= \alpha' [1 + (1 + \theta)] \leq \quad (66)$$

$$\leq 3\alpha', \quad (67)$$

and consequently

$$\begin{aligned}
\int_{\beta_1}^{\beta_2} \mathbf{E}[\Omega(h^2) - \Omega^2(h)] d\beta &\leq \int_{\beta_1}^{\beta_2} | \mathbf{E}[\Omega(h^2) - \Omega^2(h)] | d\beta = \\
&= \frac{1}{N} \int_{\beta_1}^{\beta_2} | \frac{d}{d\beta} \mathbf{E}[\Omega(h)] | d\beta \leq \\
&\leq 3 \frac{\alpha'}{N}
\end{aligned} \tag{68}$$

$$\Rightarrow \lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} \mathbf{E}[\Omega(h^2) - \Omega^2(h)] d\beta = 0 \tag{69}$$

and the proof is closed. \square

A.2 Details in Bernoulli dilution calculations

Here some technical calculations concerning the self-averaging technique on the Bernoulli diluted graph are reported.

$$\begin{aligned}
\mathbf{E}[\Omega(h_l G)] &= -\frac{1}{N} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sum_{\nu=1}^k \sigma_{i_\nu}^l \sigma_{j_\nu}^l G e^{(\beta \sum_{a=1}^s \sum_{\nu=1}^k \sigma_{i_\nu}^a \sigma_{j_\nu}^a)}}{\sum_{\{\sigma\}} e^{(\beta \sum_{a=1}^s \sum_{\nu=1}^k \sigma_{i_\nu}^a \sigma_{j_\nu}^a)}} \right] = \\
&= -\frac{1}{N} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} k \sigma_{i_0}^l \sigma_{j_0}^l G e^{(\beta \sum_{a=1}^s \sum_{\nu=1}^k \sigma_{i_\nu}^a \sigma_{j_\nu}^a)}}{\sum_{\{\sigma\}} e^{(\beta \sum_{a=1}^s \sum_{\nu=1}^k \sigma_{i_\nu}^a \sigma_{j_\nu}^a)}} \right],
\end{aligned}$$

and remembering the properties of the Bernoullian distribution (5) we can write

$$\begin{aligned}
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0}^l \sigma_{j_0}^l G \prod_{a=1}^s e^{\beta \sigma_{i_0}^a \sigma_{j_0}^a} e^{-\beta H_s}}{\sum_{\{\sigma\}} \prod_{a=1}^s e^{\beta \sigma_{i_0}^a \sigma_{j_0}^a} e^{-\beta H_s}} \right] = \\
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0}^l \sigma_{j_0}^l G \prod_{a=1}^s [\cosh \beta + \sigma_{i_0}^a \sigma_{j_0}^a \sinh \beta] e^{-\beta H_s}}{\sum_{\{\sigma\}} \prod_{a=1}^s [\cosh \beta + \sigma_{i_0}^a \sigma_{j_0}^a \sinh \beta] e^{-\beta H_s}} \right] =
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\frac{\sum_{\{\sigma\}} \sigma_{i_0}^l \sigma_{j_0}^l G \prod_{a=1}^s [1 + \sigma_{i_0}^a \sigma_{j_0}^a \theta] e^{-\beta H_s}}{\sum_{\{\sigma\}} \prod_{a=1}^s [1 + \sigma_{i_0}^a \sigma_{j_0}^a \theta] e^{-\beta H_s}} \right] = \\
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\frac{\Omega \left(\sigma_{i_0}^l \sigma_{j_0}^l G \prod_{a=1}^s [1 + \sigma_{i_0}^a \sigma_{j_0}^a \theta] \right)}{\left(1 + \omega(\sigma_{i_0}^a \sigma_{j_0}^a) \theta \right)^s} \right] =
\end{aligned}$$

let us expand both the numerator and the denominator up to the second order in θ

$$\begin{aligned}
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\Omega \left((\sigma_{i_0}^l \sigma_{j_0}^l G) \left(1 + \sum_a^{1,s} \sigma_{i_0}^a \sigma_{j_0}^a \theta + \sum_{a<b}^{1,s} \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^b \sigma_{j_0}^b \theta^2 \right) \right) \times \right. \\
&\quad \times \left. \left(1 - s\omega(\sigma_{i_0} \sigma_{j_0}) \theta + \frac{s(s+1)}{2} \omega^2(\sigma_{i_0} \sigma_{j_0}) \theta^2 \right) \right] = \\
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\Omega \left(G \sigma_{i_0}^l \sigma_{j_0}^l + G \sum_a^{1,s} \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^l \sigma_{j_0}^l \theta + G \sum_{a<b}^{1,s} \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^b \sigma_{j_0}^b \sigma_{i_0}^l \sigma_{j_0}^l \theta^2 \right) \times \right. \\
&\quad \times \left. \left(1 - s\omega(\sigma_{i_0}^a \sigma_{j_0}^a) \theta + \frac{s(s+1)}{2} \omega^2(\sigma_{i_0} \sigma_{j_0}) \theta^2 \right) \right] = \\
&= -\frac{\alpha M}{N^2} \mathbf{E} \left[\Omega(G \sigma_{i_0}^l \sigma_{j_0}^l) + \theta \left(\sum_a^{1,s} \Omega(G \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^l \sigma_{j_0}^l) - s\Omega(G \sigma_{i_0}^l \sigma_{j_0}^l) \omega(\sigma_{i_0} \sigma_{j_0}) \right) + \right. \\
&\quad + \theta^2 \left(\sum_{a<b}^{1,s} \Omega(G \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^b \sigma_{j_0}^b \sigma_{i_0}^l \sigma_{j_0}^l) - s \sum_a^{1,s} \Omega(G \sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^l \sigma_{j_0}^l) \omega(\sigma_{i_0} \sigma_{j_0}) + \right. \\
&\quad \quad \left. + \frac{s(s+1)}{2} \Omega(G \sigma_{i_0}^l \sigma_{j_0}^l) \omega^2(\sigma_{i_0}^a \sigma_{j_0}^a) \right) \Big] = \\
&= -\frac{\alpha M}{N^2} \left[\langle G m_l^2 \rangle + \theta \left(\sum_{a=1}^s \langle G q_{a,l}^2 \rangle - s \langle G q_{l,s+1}^2 \rangle \right) + \right. \\
&\quad \left. + \theta^2 \left(\sum_{a<b}^{1,s} \langle G q_{l,a,b}^2 \rangle - s \sum_a^{1,s} \langle G q_{l,a,s+1}^2 \rangle + \frac{s(s+1)}{2} \langle G q_{l,s+1,s+2}^2 \rangle \right) \right].
\end{aligned}$$

While the other term $\mathbf{E}[\Omega(h_l)\Omega(G)]$ can be worked out as follows:

$$\begin{aligned}
& \mathbf{E}[\Omega(h_l)\Omega(G)] = \tag{70} \\
&= -\frac{1}{N}\mathbf{E}\left[\frac{\sum_{\{\sigma\}}\sum_{\nu=1}^k\sigma_{i_\nu}^l\sigma_{j_\nu}^l G e^{\beta\sum_{\nu=1}^k\sigma_{i_\nu}^l\sigma_{j_\nu}^l} e^{(\beta\sum_{a=1}^s\sum_{\nu=1}^k\sigma_{i_\nu}^a\sigma_{j_\nu}^a)}}{\sum_{\{\sigma\}}e^{\beta\sum_{\nu=1}^k\sigma_{i_\nu}^l\sigma_{j_\nu}^l} e^{(\beta\sum_{a=1}^s\sum_{\nu=1}^k\sigma_{i_\nu}^a\sigma_{j_\nu}^a)}}\right] = \\
&= -\frac{1}{N}\mathbf{E}\left[\frac{\sum_{\{\sigma\}}k\sigma_{i_0}^l\sigma_{j_0}^l G e^{\beta\sum_{\nu=1}^k\sigma_{i_\nu}^l\sigma_{j_\nu}^l} e^{(\beta\sum_{a=1}^s\sum_{\nu=1}^k\sigma_{i_\nu}^a\sigma_{j_\nu}^a)}}{\sum_{\{\sigma\}}e^{\beta\sum_{\nu=1}^k\sigma_{i_\nu}^l\sigma_{j_\nu}^l} e^{(\beta\sum_{a=1}^s\sum_{\nu=1}^k\sigma_{i_\nu}^a\sigma_{j_\nu}^a)}}\right] = \\
&= -\frac{\alpha M}{N^2}\mathbf{E}\left[\frac{\sum_{\{\sigma\}}\sigma_{i_0}^l\sigma_{j_0}^l G e^{\beta\sigma_{i_0}^l\sigma_{j_0}^l} [\prod_{a=1}^s e^{\beta\sigma_{i_0}^a\sigma_{j_0}^a}] e^{-\beta H_{s+1}}}{\sum_{\{\sigma\}}e^{\beta\sigma_{i_0}^l\sigma_{j_0}^l} [\prod_{a=1}^s e^{\beta\sigma_{i_0}^a\sigma_{j_0}^a}] e^{-\beta H_{s+1}}}\right] = \\
&= -\frac{\alpha M}{N^2}\mathbf{E}\left[\frac{\Omega\left(\sigma_{i_0}^l\sigma_{j_0}^l G(1+\sigma_{i_0}^l\sigma_{j_0}^l\theta)[\prod_{a=1}^s(1+\sigma_{i_0}^a\sigma_{j_0}^a\theta)]\right)}{(1+\omega(\sigma_{i_0}\sigma_{j_0})\theta)^{s+1}}\right] = \\
&= -\frac{\alpha M}{N^2}\mathbf{E}\left[\Omega\left((\sigma_{i_0}^l\sigma_{j_0}^l G + G\theta)(1+\sum_a^{1,s}\sigma_{i_0}^a\sigma_{j_0}^a\theta + \sum_{a<b}^{1,s}\sigma_{i_0}^a\sigma_{j_0}^a\sigma_{i_0}^b\sigma_{j_0}^b\theta^2)\right) \times \right. \\
&\quad \times \left. \left(1-(s+1)\omega(\sigma_{i_0}\sigma_{j_0})\theta + \frac{(s+1)(s+2)}{2}\omega^2(\sigma_{i_0}\sigma_{j_0})\theta^2\right)\right] = \\
&= -\frac{\alpha M}{N^2}\mathbf{E}\left[\left(\Omega(G\sigma_{i_0}^l\sigma_{j_0}^l) + \theta\left(\Omega(G) + \sum_a^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a\sigma_{i_0}^l\sigma_{j_0}^l)\right) + \right. \right. \\
&\quad \left. \left. + \theta^2\left(\sum_a^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a) + \sum_{a<b}^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a\sigma_{i_0}^b\sigma_{j_0}^b\sigma_{i_0}^l\sigma_{j_0}^l)\right)\right) \times \right. \\
&\quad \times \left. \left(1-(s+1)\omega(\sigma_{i_0}\sigma_{j_0})\theta + \frac{(s+1)(s+2)}{2}\omega^2(\sigma_{i_0}\sigma_{j_0})\theta^2\right)\right] = \\
&= -\frac{\alpha M}{N^2}\mathbf{E}\left[\Omega(G\sigma_{i_0}^l\sigma_{j_0}^l) + \theta\left(\Omega(G) + \sum_a^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a\sigma_{i_0}^l\sigma_{j_0}^l) \right. \right. \\
&\quad \left. \left. - (s+1)\Omega(G\sigma_{i_0}^l\sigma_{j_0}^l)\omega(\sigma_{i_0}\sigma_{j_0})\right) + \right. \\
&\quad \left. + \theta^2\left(\sum_a^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a) + \sum_{a<b}^{1,s}\Omega(G\sigma_{i_0}^a\sigma_{j_0}^a\sigma_{i_0}^b\sigma_{j_0}^b\sigma_{i_0}^l\sigma_{j_0}^l) \right. \right. \\
&\quad \left. \left. - (s+1)\omega(\sigma_{i_0}\sigma_{j_0})\Omega(G\sigma_{i_0}^l\sigma_{j_0}^l) + \frac{(s+1)(s+2)}{2}\omega^2(\sigma_{i_0}\sigma_{j_0})\Omega(G)\right)\right] = \tag{71}
\end{aligned}$$

$$\begin{aligned}
& + \theta^2 \left(\sum_a^{1,s} \Omega(G\sigma_{i_0}^a \sigma_{j_0}^a) + \sum_{a < b}^{1,s} \Omega(G\sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^b \sigma_{j_0}^b \sigma_{i_0}^l \sigma_{j_0}^l) \right. \\
& - (s+1)\Omega(G)\omega(\sigma_{i_0} \sigma_{j_0}) + \\
& - (s+1) \sum_a^{1,s} \Omega(G\sigma_{i_0}^a \sigma_{j_0}^a \sigma_{i_0}^l \sigma_{j_0}^l) \omega(\sigma_{i_0} \sigma_{j_0}) \\
& \left. + \frac{(s+1)(s+2)}{2} \Omega(G\sigma_{i_0}^l \sigma_{j_0}^l) \omega^2(\sigma_{i_0}^a \sigma_{j_0}^a) \right) = \\
& = -\frac{\alpha M}{N^2} \left[\langle Gm_l^2 \rangle + \theta \left(\langle G \rangle + \sum_{a=1}^s \langle Gq_{a,l}^2 \rangle - (s+1) \langle Gq_{l,s+1}^2 \rangle \right) + \right. \\
& + \theta^2 \left(\sum_a^{1,s} \langle Gm_a^2 \rangle - (s+1) \langle Gm_l^2 \rangle + \sum_{a < b}^{1,s} \langle Gq_{l,a,b}^2 \rangle + \right. \\
& \left. - (s+1) \sum_a^{1,s} \langle Gq_{l,a,s+1}^2 \rangle + \frac{(s+1)(s+2)}{2} \langle Gq_{l,s+1,s+2}^2 \rangle \right) \Big]
\end{aligned}$$

A.3 Poisson identities via the self-averaging technique

For the sake of completeness we report also the constraints in the Poisson diluted model obtained by using the first method:

$$\begin{aligned}
f_G(\alpha, \beta) &= \alpha \left[\left(\sum_{l=1}^s \langle Gm_l^2 \rangle - s \langle Gm_{s+1}^2 \rangle \right) (1 - \theta^2) + \right. \\
& + 2\theta \left(\sum_{a < l}^{1,s} \langle Gq_{al}^2 \rangle - s \sum_l^{1,s} \langle Gq_{l,s+1}^2 \rangle + \frac{s(s+1)}{2} \langle Gq_{s+1,s+2}^2 \rangle \right) + \\
& + 3\theta^2 \left(\sum_{l < a < b}^{1,s} \langle Gq_{l,a,b}^2 \rangle - s \sum_{l < a}^{1,s} \langle Gq_{l,a,s+1}^2 \rangle + \frac{s(s+1)}{2} \sum_l^{1,s} \langle Gq_{l,s+1,s+2}^2 \rangle + \right. \\
& \left. - \frac{s(s+1)(s+2)}{3!} \langle Gq_{s+1,s+2,s+3}^2 \rangle \right) + O(\theta^3) \Big], \tag{72}
\end{aligned}$$

by which, choosing as a trial function m^2 we have

$$\begin{aligned} f_{m^2}^P(\alpha, \beta) &= \alpha \left[\left(\langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle \right) (1 - \theta^2) + \right. \\ &\quad \left. - 2\theta \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle \right) + \right. \\ &\quad \left. + 3\theta^2 \left(\langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle \right) + O(\theta^3) \right], \end{aligned}$$

from which, changing the Jacobian $d\theta = (1 - \theta^2)d\beta$, we get

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} |f_{m^2}^P(\alpha, \beta)| d\beta &= \alpha \int_{\theta_1}^{\theta_2} d\theta \left[\left(\langle m_1^4 \rangle - \langle m_1^2 m_2^2 \rangle \right) + \right. \\ &\quad - 2 \frac{\theta}{(1 - \theta^2)} \left(\langle m_1^2 q_{12}^2 \rangle - \langle m_1^2 q_{23}^2 \rangle \right) + \\ &\quad + 3 \frac{\theta^2}{(1 - \theta^2)} \left(\langle m_1^2 q_{123}^2 \rangle - \langle m_1^2 q_{234}^2 \rangle \right) \\ &\quad \left. + O(\theta^3) \right] = 0. \end{aligned} \quad (73)$$

If we set $G = q_{12}^2$ as the trial function $f_G^P(\alpha, \beta)$ becomes

$$\begin{aligned} f_{q^2}^P(\alpha, \beta) &= \alpha \left[\left(2 \langle m_1^2 q_{12}^2 \rangle - 2 \langle m_3^2 q_{12}^2 \rangle \right) (1 - \theta^2) + \right. \\ &\quad + 2\theta \left(\langle q_{12}^4 \rangle - 4 \langle q_{12}^2 q_{23}^2 \rangle + 3 \langle q_{12}^2 q_{34}^2 \rangle \right) + \\ &\quad \left. - 6\theta^2 \left(\langle q_{12}^2 q_{123}^2 \rangle - 3 \langle q_{12}^2 q_{234}^2 \rangle + 2 \langle q_{12}^2 q_{345}^2 \rangle \right) + O(\theta^3) \right]. \end{aligned}$$

Again

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\beta_1}^{\beta_2} |f_{q^2}^P(\alpha, \beta)| d\beta &= 2\alpha \int_{\theta_1}^{\theta_2} d\theta \left[\left(\langle m_1^2 q_{12}^2 \rangle - \langle m_3^2 q_{12}^2 \rangle \right) + \right. \\ &\quad + \frac{\theta}{(1 - \theta^2)} \left(\langle q_{12}^4 \rangle - 4 \langle q_{12}^2 q_{23}^2 \rangle + 3 \langle q_{12}^2 q_{34}^2 \rangle \right) + \\ &\quad - 3 \frac{\theta^2}{(1 - \theta^2)} \left(\langle q_{12}^2 q_{123}^2 \rangle - 3 \langle q_{12}^2 q_{234}^2 \rangle + 2 \langle q_{12}^2 q_{345}^2 \rangle \right) \\ &\quad \left. + O(\theta^3) \right] = 0, \end{aligned} \quad (74)$$

References

- [1] E. Agliari, A. Barra, F. Camboni, *Criticality in diluted ferromagnets*, J. Stat. Mech. P1003 (2008).
- [2] M. Aizenman, P. Contucci, *On the stability of the quenched state in mean field spin glass models*, J. Stat. Phys. **92**, 765-783 (1998).
- [3] R. Albert, A. L. Barabasi *Statistical mechanics of complex networks*, Reviews of Modern Physics **74**, 47-97 (2002).
- [4] A. Barra, *The mean field Ising model through interpolating techniques*, J. Stat. Phys. **145**, 234-261, (2008).
- [5] A. Barra *Notes of ferromagnetic Pspin and REM*, Math. Meth. Appl. Sci. 10.1002/mma.1065, Wiley (2008).
- [6] A. Barra, *Irreducible free energy expansion and overlap locking in mean field spin glasses*, J. Stat. Phys. **123**, 601-614 (2006).
- [7] A. Barra, L. DeSanctis *Stability properties and probability distribution of multi-overlaps in dilute spin glasses*, J. Stat. Mech. P08025 (2007).
- [8] A. Barra, F. Guerra, *About the ergodicity in Hopfield analogical neural networks*, J. Math. Phys. **49** 125217 (2008).
- [9] A. Barra, F. Guerra, *Order parameters and their locking in analogical neural networks*, "Percorsi Incrociati", Dedicated Volume, Collana d'Ateneo, University of Salerno (2008).
- [10] A. Barrat, M. Weight, *On the properties of small world network models*, Eur. Phys. Journ. B **13**-3 (2000).
- [11] A. Bianchi, P. Contucci, A. Knauf, *Stochastically Stable Quenched Measures*, Journal of Statistical Physics, **117**, 831-844, (2004).
- [12] M. Buchanan, *Nexus: Small Worlds and the Groundbreaking Theory of Networks*. Norton, W. W. Company, Inc. (2003).

- [13] F. Chung, L. Lu, *Complex Graphs and Networks*, Americ. Math. Soc. Publ. (2006).
- [14] P. Contucci, C. Giardiná, *The Ghirlanda-Guerra identities*, J. Stat. Phys., **126**, N. 4/5, 917-931, (2007).
- [15] P. Contucci, C. Giardiná, I. Nishimori, *Spin Glass Identities and the Nishimori Line*, to appear in "Progress in Probability", (2008).
- [16] P. Contucci, J. Lebowitz, *Correlation Inequalities for Spin Glasses*, Annales Henri Poincare, **8**, 1461-1467, (2007).
- [17] P. Contucci, M. Degli Esposti, C. Giardiná, S. Graffi, *Thermodynamical Limit for Correlated Gaussian Random Energy Models*, Commun. in Math. Phys. **236**, 55-63, (2003).
- [18] A. Dembo, A. Montanari, *Ising models on locally tree-like graphs*, arXiv:0804.4726 (2008).
- [19] R.S. Ellis, *Large deviations and statistical mechanics*, Springer, New York (1985).
- [20] S. Ghirlanda, F. Guerra, *General properties of overlap distributions in disordered spin systems. Towards Parisi ultrametricity*, J. Phys. A, **31**, 9149-9155, (1998).
- [21] F. Guerra, *About the overlap distribution in mean field spin glass models*, Int. Jou. Mod. Phys. B **10**, 1675-1684, (1996).
- [22] F. Guerra, L. De Sanctis, *Mean field dilute ferromagnet I. High temperature and zero temperature behavior*, J. Stat. Phys. **129**, 231, (2008).
- [23] F. Guerra, F. L. Toninelli, *The Thermodynamic Limit in Mean Field Spin Glass Models*, Comm. Math. Phys. **230**, 71-79, (2002).
- [24] F. Guerra, F. L. Toninelli, *The high temperature region of the Viana-Bray diluted spin glass model*, J. Stat. Phys. **115**, (2004).
- [25] M. Mezard, G. Parisi, R. Zecchina, *Analytic and Algorithmic Solution of Random Satisfiability Problems*, Science **297**, 812 (2002).

- [26] R. Monasson, S.Kirkpatrick, B.Selman, L.Troyansky, R. Zecchina, *Determining computational complexity from characteristic phase transitions*, Nature **400**, 133 (1999).
- [27] M. Newman, D. Watts, A.-L. Barabasi *The Structure and Dynamics of Networks*, Princeton University Press, (2006).
- [28] T. Nikoletopolous, A.C.C. Coolen, I. Perez Castillo, N.S.Skantos, J.P.L. Hatchett, B. Wemmenthove, *Replicated transfer matrix analysis of Ising spin models on small world lattices*, J. Phys. A, 6455-6475 (2004).
- [29] L. Pastur, M. Shcherbina, *The absence of self-averaging of the order parameter in the Sherrington-Kirkpatrick model*, J. Stat. Phys. **62**, 1-19, (1991).
- [30] D.J. Watts, S.H. Strogatz, *Collective dynamics of small world networks*, Nature **393**, 6684 (1998)